

Sunlight Sonata: Contrasting the Roles of Solar Ultraviolet Radiation and Temperature on the Calling Behavior of a Diurnal Forest-specialist Frog

Álvaro A. Mainardi¹, Vinícius M. Caldart², Mauricio Beux dos Santos³, Nelson J. Schuch⁴,
and André P. Schuch^{1,5,*}

¹Programa de Pós-Graduação em Biodiversidade Animal, Universidade Federal de Santa Maria, Building 17, Room 1140-D, Roraima Avenue, Camobi, Santa Maria, RS, 97105-900, Brazil. E-mail: alvaromainardi@biodos.com.br (Mainardi)

²Área das Ciências da Vida e Saúde, Universidade do Oeste de Santa Catarina, Rua José Firmo Bernardi, 1591, 89600-000, Joaçaba, Santa Catarina, Brazil. E-mail: viniciuscaldart@gmail.com (Caldart)

³BioDos Soluções em Dosimetria Ltda, Roraima Avenue, 1000, Camobi, Santa Maria, RS, 97105-340, Brazil. E-mail: mbeuxs@gmail.com (Beux dos Santos)

⁴Coordenação Espacial do Sul (COESU/INPE-MCTI), Universidade Federal de Santa Maria, Santa Maria, RS, 97105-970, Brazil. E-mail: njschuch@gmail.com (Schuch)

⁵Department of Biochemistry and Molecular Biology, Universidade Federal de Santa Maria, Building 74E, Room 16, Roraima Avenue, Camobi, Santa Maria, RS, 97105-900, Brazil. *Correspondence: E-mail: andre.schuch@ufsm.br (Schuch)

Received 24 May 2025 / Accepted 7 March 2026 / Published 2 July 2026
Communicated by Tzu-Hao Lin

Diurnal animals are frequently exposed to solar ultraviolet radiation (UV) during their daily activities, which can lead to DNA damage and other physiological challenges. To counteract these effects, they have evolved complex DNA repair mechanisms. The high energetic cost of DNA repair creates a trade-off with behavior, potentially constraining energetically demanding activities like anuran calling under harmful UV exposure. We tested hypotheses on how natural heterogeneity in the abiotic environment – particularly in the UV radiation – influences the calling behavior of *Crossodactylus schmidtii*, a diurnal forest-specialist frog found in preserved neotropical streams. To gather data on male calling behavior, we used passive acoustic monitoring of a well-studied population, using a set of recorders and data loggers that measured short- and long-term cumulative doses of UVA and UVB, air temperature, light intensity and cloud cover during the austral summer. To investigate associations between the abiotic variables and proxies of calling behavior (occurrence, number, and duration of calls), we used a model selection approach based on generalized linear mixed models followed by model averaging of the best-supported models. We found that: 1) the probability of recording males in calling behavior decreased with increasing mean air temperature and cumulative UVB dose but increased with short-term (15-minute) UVB exposure; 2) the number of calls emitted by males decreased with increasing mean air temperature but increased with cumulative UVB dose; and 3) call duration decreased with increasing mean air temperature but increased with cumulative UVB dose at calling sites. Short- and long-term doses of UVA, on the other hand, did not influence any proxy of calling behavior. Our findings show that higher temperatures in months of peak calling activity consistently reduce calling behavior, while UVB exposure has more complex effects. Together, these results underscore the need to consider both thermal and ultraviolet components of sunlight when assessing how anurans respond behaviorally to abiotic heterogeneity under ongoing environmental and climatic changes.

Keywords: Air Temperature, *Crossodactylus*, Environmental Heterogeneity, Hylodidae, Reproductive Behavior, Solar Ultraviolet Radiation

Citation: Mainardi AA, Caldart VM, Beux dos Santos M, Schuch NJ, Schuch AP. 2026. Sunlight sonata: contrasting the roles of solar ultraviolet radiation and temperature on the calling behavior of a diurnal forest-specialist frog. Zool Stud 65:22. doi:10.6620/ZS.2026.65-22.

BACKGROUND

Diurnal animals have evolved under exposure to solar ultraviolet radiation (UV). The two spectra of UV radiation that reach the Earth's surface are UVA, which has longer and less energetic wavelengths (315–400 nm), and UVB, of shorter and more energetic wavelengths (280–315 nm) (Frederick et al. 1989). UVB is more dangerous to organisms because of a higher absorbance by the DNA molecule, while UVA has a lower absorbance and requires a longer exposure time to generate similar damage relative to UVB (Hamzavi and Lui 2005; Markovitsi et al. 2010; Schuch et al. 2017). The consequences of exposure to UV on organisms—which include genetic, physiological, and behavioral effects—are therefore wavelength-dependent and dose-dependent (reviewed in Londero et al. 2019). For example, the negative effects of UVB on echinoderm embryo development increase with exposure time (Lesser and Barry 2003), fish mortality increases with accumulated UVB irradiance, and prolonged exposure to UVA causes skin lesions (Zagarese and Williamson 2001; Kouwenberg et al. 1999). However, animals such as anurans and other vertebrates have evolved adaptive responses to UV radiation exposure (reviewed in Cockell and Blaustein 2001), including biochemical mechanisms to repair UV-induced DNA damage (Lamare et al. 2006; Dong et al. 2007; Kienzler et al. 2013; Cortat et al. 2013) and the capacity to benefit from low doses of UVB light for metabolic regulation, such as in vitamin D synthesis (Neer 1975; Xue and Fleet 2009; Hohman et al. 2011).

Animals typically concentrate their active periods at specific times of the day as strategies to cope with the environment's temporal variation in abiotic factors (Daan 1981). Nonetheless, diurnal animals that remain active for prolonged times, or when UV radiation is most intense, may be exposed to UV radiation that exceeds their physiological optimum and induces DNA damage. Photoproducts such as Cyclobutane pyrimidine dimers and [6-4] pyrimidine-pyrimidone can disrupt homeostasis by causing mutations and impairing cellular functions (Schuch and Menck 2010). In fact, even small doses of UV radiation are genotoxic at the cellular level and can cause DNA damage (Schuch et al. 2017; Yagura et al. 2017; Londero et al. 2019). In vertebrates, the DNA repair mechanism counteracts this damage, but because the mechanism is physiologically costly, it can limit energy allocation in certain behaviors (García-Huidobro et al. 2017; Pulgar et al. 2017). For example, an energetic trade-off between behavior and physiology due to exposure to harmful UV radiation can be discerned through a focused examination of anuran calling behavior.

Amphibian studies provide insightful findings on the influence of UV exposure on animal behavior, as these organisms lack protective skin appendages, and their eggs, lacking shells, are highly susceptible to environmental variations (Blaustein and Belden 2003). For example, adult frogs exposed to a minor increase in daily UV dose reduce their investment in locomotion and in selection and use of calling sites (Blaustein et al. 2000; Kats et al. 2012), whereas tadpoles are more susceptible to predation and less capable of foraging under UV exposure (Kats et al. 2000; Londero et al. 2017; Martins et al. 2021). Forest-specialist diurnal amphibians are expected to be more vulnerable to UV effects than open-habitat species due to their lower concentration of melanin in the skin (Blaustein and Belden 2003), which acts as a protective barrier for the UV spectrum (reviewed in Brenner and Hearing 2008). However, studies examining how UV heterogeneity influences the behavior of diurnal amphibians in forest streams are scarce and mostly limited to laboratory conditions (see Kats et al. 2012; Schuch et al. 2015). This gap is particularly striking because most amphibians are nocturnal and thus rarely exposed to solar UV radiation, making diurnal lineages especially valuable for understanding behavioral responses to UV.

Here we aimed to answer the following question: how does the abiotic environment modulate male frogs' calling behavior in forest streams? We used *Crossodactylus schmidtii*, a diurnal forest-specialist frog found in preserved neotropical streams, as a study species. Unlike most amphibians, which avoid sunlight, *C. schmidtii* is active throughout the day under natural and varying levels of UV exposure, making it an exceptional model to assess how solar radiation influences animal behavior in the field. The ecology and behavior of *C. schmidtii* offer a unique opportunity to investigate, in natural settings, how heterogeneity in the abiotic environment modulates the behavior of diurnal amphibians in forest streams. Males of *C. schmidtii* compete for and defend territories in the stream bed, emitting acoustic, visual and multimodal signals during contests (Caldart et al. 2014 2022), from sunrise to sunset, throughout the year (Caldart et al. 2016).

In this study, we deployed data loggers in a stream to monitor ultraviolet radiation, air temperature, and luminosity for extended periods, while audio equipment simultaneously recorded the calling behavior of males of *C. schmidtii*. We then tested two hypotheses on how abiotic factors correlate with calling behavior: (1) high UV doses at male calling sites are associated with a lower probability of recording males in calling behavior; and (2) the investment of males in calling behavior (*i.e.*, in number and duration of calls) decreases with increasing UV incidence in calling sites. Overall, we

expect UVB to have stronger impacts on male calling behavior than UVA, as it is more readily absorbed by DNA, causes faster damage, and demands more energy for repair.

MATERIALS AND METHODS

Study Area

The study was carried out at Turvo State Park, located in the municipality of Derrubadas, northwest of Rio Grande do Sul, Brazil (27°13'57.61"S, 53°51'4.82"W), on the border with Argentina. The park vegetation is semi-deciduous Atlantic Forest and covers an area of 17,491 ha (SEMA 2005). The local climate is subtropical humid, characterized by temperatures above 22°C in January and minimum temperature close to -3°C in July (Alvares et al. 2013). The annual mean precipitation is 1,787 mm, with rainfall evenly distributed throughout the year. Summer occurs in December–March and winter in June–September (Maluf 2000).

Data Sampling

The data were collected from December 2019 to January 2020, the warmer months that correspond to the austral summer, in which there is a high proportion of sexually mature adults of *Crossodactylus schmidtii* in the study area and male calling behavior peaks (Caldart et al. 2016). As individuals of this species are not regularly distributed between and within streams due to spatial variation in the availability of calling and breeding sites, we visited streams beforehand to select a stream having active males in high density to conduct the study (see Caldart et al. 2016). We chose a first-order stream that presents an extensive canopy cover (27°14'01.7"S, 53°57'52.9"W). Along the stream we established three fixed sites with confirmed occurrence of males, in which we deployed data loggers and conducted passive acoustic monitoring (Fig. 1). The monitoring sites were separated from each other by at least 50 m, which is a sufficient distance to assure acoustic independence in a forest-lotic habitat (Caldart et al. 2016; Statistical Analysis section, Fig. 1B).

At each site we used passive acoustic monitoring to record the vocalizations emitted by males between 07:00 and 18:00 h, for 19 days, using a pair of Sony ICD-PX312 digital recorders (following Acevedo and Villanueva-Rivera 2006; Blumstein et al. 2011) positioned in opposite stream banks and at 40 cm from the streambed (*i.e.*, the calling sites, Fig. 1C). The audio recordings were combined into a stereo file using

PluralEyes 4 software (Red Giant). The first 5 mins at every 15-minute time interval (time hereafter) were analyzed using the software Audition (Adobe Audition CC 2018). From the stereo files we then obtained the following data for each time interval analyzed, at each site and day of monitoring: 1) either the occurrence or the absence of individuals in calling behavior, 2) the total number of calls emitted by all males, and 3) the mean duration of the calls emitted by all males. In the analysis of the recordings, we considered only advertisement calls because it is a call frequently emitted by males throughout the day (Caldart et al. 2016), is higher pitched, louder, and has higher inter-individual temporal segregation in emission than other calls known for the species (Caldart et al. 2011).

Simultaneously with the acoustic monitoring at each of the three fixed sites we measured the incidence of solar UVB (kJ/m²) and UVA (kJ/m²) radiation, luminosity, and air temperature with data loggers we built. Each data logger has a battery and a secure digital card for data recording that allows monitoring for at least 45 days without recharging or changing the battery. The device is registered and under confidentiality in the Instituto Nacional da Propriedade Industrial (INPI, Brazil) under the number BR 10 2021 005008 0 (see validation test in SM. 1). We obtained the following data from the data loggers for each 15 mins time interval, at each site and day of monitoring: the 15 mins ultraviolet dose of UVA and UVB, and the cumulative (from 07:00 h) ultraviolet dose of UVA and UVB. The daily dose in a clear sky day for UVB in the fixed sampled sites ranged from 0.02 kJ/m² to 0.45 kJ/m², while for UVA ranged from 3.36 kJ/m² to 104.59 kJ/m². This dose is equivalent to 2.69% and 6.06% of the daily dose in a clear sky day out of the forest, in the same period and latitude (see SM. 2 for more details on the profiles of UV incidence in the fixed samples sites). Because the calling behavior of diurnal frogs, including *C. schmidtii*, is often correlated with abiotic factors including luminosity and temperature (Caldart et al. 2016; Hatano et al. 2002; Van Sluys and Guido-Castro 2011), we also obtained from the data loggers the mean air temperature and the mean light intensity for each 15 min time interval at each site and day of monitoring.

Finally, because there is a relationship between cloud cover and UV radiation incidence, we obtained the cloud cover data from the Instituto Nacional de Pesquisas Espaciais (INPE, Brazil) for the same period as for the previous variables. These measures are based on an index of cloud coverage derived from the corrected reflectance values obtained in satellite imagery (Souza et al. 2013).

Statistical Analyses

To test the predictions of our hypotheses we used as response variables the 1) occurrence of calling behavior, the 2) number of calls emitted, and the 3) mean duration of calls. The first variable was used as a proxy of calling behavior (*i.e.*, whether males are acoustically active or not), whereas the other two variables were used as proxies of calling investment (*i.e.*, how much males invest in call frequency and duration). We used data from the period of 07:00 to 18:00 h (Fig. 1A), as *C. schmidtii* has a diurnal activity spanning from sunrise to sunset (Caldart et al. 2016). For each response variable we used a four-step analysis protocol that was the most suitable considering the intrinsic characteristics of our data, which contained autocorrelated observations collected through a passive

acoustic monitoring method, several covariates, and random terms of different structures.

In the first step we fitted a global (full) Generalized Linear Mixed Model (GLMM) to the original dataset to establish a baseline model that captures the relationship between the response variable and the predictor variables. We considered different combinations of response variables, error distributions and link functions (SM. 3) in the package *glmmTMB* v. 1.1.8 (Brooks et al. 2017). To control the variation in the response variables among sites, days, and time, we fitted these variables as random factors. The random factor time was fitted as a first order autoregressive term (AR1), that relates the observation at the current time, t , to the observation at the previous time, $t-1$ (Larsen et al. 2021). The model also accounted for the fact that observations close in time within the same date group




Fig. 1. Schematic representation of data collection. (A) Purple arrows and dashed lines denote the incidence of ultraviolet radiation (UVB and UVA) varying according to the solar angle and canopy cover from 07:00 to 18:00 h, which is the whole time interval between sunrise (up arrow) and sunset (down arrow). (B) Distribution of the three fixed points of acoustic monitoring (red-filled circles) along the stream (blue dashed line), showing the minimum distance between them. (C) Example of the centralized positioning of the datalogger and opposed placement of audio recording equipment in the fixed sampling points.

may be more correlated than observations further apart in time or in different date groups (*i.e.*, $\text{ar1}(\text{time} + 0 \mid \text{date})$ in R notation).

In the second step we performed bootstrap resampling to assess the stability and variability of the global model, using the package *lme4* v. 1.1.37 (Bates et al. 2015). Random resampling with replacement of the original model estimates were computed and repeated through 1000 iterations, and the stability of the model estimates were assessed comparing the curves shapes, median values and quantile ranges of original and bootstrap estimates in Kernel density estimation plots (SMs. 4–6). Bootstrap resampling helps in understanding how stable the estimates from the global model are and provides a distribution of parameter estimates that reflects the variability in the data (Davison and Hinkley 1997).

In the third step we performed a model selection approach (model dredging) upon the global model to explore alternative models that may better explain the variability in the data. Finally, in the fourth step we used a model averaging procedure to obtain a consolidated estimate by combining the subset of models with $\Delta\text{AIC} < 2$ obtained in step 3, as implemented in the *MuMIn* package v.1.48.11 (Bartón 2025). Model averaging considers the uncertainty associated with model selection by assigning weights to each model's predictions and combining them to obtain a final estimate, providing a more robust estimate by considering multiple candidate models (Symonds and Moussalli 2011). To ensure comparability across models with different response distributions (binomial, gamma, and gaussian) and correlation structures, we quantified model fit using the pseudo- R^2 metric, computed as the squared Pearson correlation between observed and predicted values (Nakagawa and Schielzeth 2013).

In the model designed to examine the hypothesis that high UV doses at male calling sites correlate with a lower probability of males engaging in calling behavior, we introduced the response variable occurrence of calling behavior. This variable adopts binary values (0 or 1), where 0 represents the absence of male vocalizations during a specified time interval, and 1 indicates the presence of vocalizations, regardless of the number of males or the frequency of calls emitted. The predictor variables were the time correspondent values of cloud cover, mean luminosity, mean air temperature, 15-minute dose of UVA, 15-minute cumulative dose of UVB, cumulative dose of UVA (from 07:00 h to the observation at the current time, t) and cumulative dose of UVB (from 07:00 h to t).

In the models built to test the hypothesis that the calling investment decreases with increasing UV incidence at calling sites, we entered the number of calls

emitted and the duration of calls as response variables. The number of calls emitted, and the duration of calls, represent the total and the mean observed values within each time interval, respectively. Predictor variables were the same as in the first model. These models included only non-zero values in the response variable (*i.e.*, only active males) as we were interested in exploring further the association between UV and the calling investment of individuals that remained acoustically active. For all models, the cloud cover was the only predictor entered as an interaction term with 15-minute dose of UVA, 15-minute dose of UVB, mean light intensity, and mean air temperature, because cloud cover can affect these variables in different ways. For example, cloud cover can either act as a major barrier to UV after the ozone layer or can diffuse the UV in the environment when there is a greater number of water particles in the air (Frederick et al. 1989).

Finally, to evaluate whether the number of calls and the mean call duration differed among periods of the day (*i.e.*, morning hours, midday hours and afternoon hours), we fitted generalized linear mixed-effects models (GLMMs) using the *glmmTMB* v. 1.1.8 (Brooks et al. 2017). In both models, the period of the day (either 07:00–10:45 h, 11:00–13:45 h or 14:00–17:45 h) was included as a categorical fixed effect. To account for the non-independence of observations taken at the same sites and on the same dates, we included random intercepts for site and date in both models. As in the previous models, the random factor time was fitted as a first order autoregressive term (AR1), that relates the observation at the current time, t , to the observation at the previous time, $t-1$ (see Larsen et al. 2021), and the model accounted for correlation between observations close in time within the same date (*i.e.*, $\text{ar1}(\text{time} + 0 \mid \text{date})$ in R notation). Estimated marginal means (EMMs) and pairwise contrasts among periods of the day were obtained using the *emmeans* package v 1.11.2.8 (Lenth and Piaskowski 2025) with Tukey's correction for multiple comparisons.

Rainfall and air humidity were not included in the models because previous work on the same population showed no effect of air humidity on daily calling activity and demonstrated that accumulated rainfall influences calling behavior only at monthly, but not daily, temporal scales (Caldart et al. 2016). Furthermore, *C. schmidtii* males occupy permanent lotic environments with consistently high ambient humidity and, as a result, males do not experience meaningful humidity gradients during daily calling activity (Caldart et al. 2016). In all models, the continuous predictors were standardized and centered to zero to produce comparable effect sizes (Schielzeth 2010). When fitting a model, we used the variance inflation factor (VIF) to analyze the magnitude

of multicollinearity of model terms using the package Performance (Lüdtke et al. 2021), and kept in models only the predictors that presented $VIF < 4$ with other predictors (a VIF value > 5 indicates a problematic amount of collinearity of a given predictor with other predictors; James et al. 2013). Finally, after fitting a model we tested its goodness of fit, the significance of the dispersion parameter (to check for under- or over-dispersion), and the presence or absence of zero-inflation, when applicable, using package *DHARMa* version 0.4.7 (Hartig 2022). Diagnostic plots and diagnostic tests for all models are presented in SMs. 4–6. All statistical analyses were performed in R version 4.1.2 (R Core Team 2022).

RESULTS

The 19 days of passive acoustic monitoring totaled 484 hours and 15 mins of recordings, of which 193 hours and 45 mins (40.1% of total recording time) included males exhibiting calling activity. The occurrence of calling activity was more common during the morning hours (53.1% of total morning time intervals analyzed), intermediate at midday hours (42.6%), and lower in the afternoon hours (25%; Fig. 2A).

The best-supported models ($\Delta AIC_c < 2$) to explain the occurrence of male calling behavior contained three models with ΔAIC_c weights (w_i) varying from 0.52 to 0.24 (Table 1). The best-ranked model included air temperature, 15-minute dose of UVB, and cumulative dose of UVB as predictors ($\Delta AIC_c w_i = 0.52$). The other

two models included air temperature, 15-minute dose of UVB, cumulative dose of UVB, and cumulative dose of UVA ($\Delta AIC_c w_i = 0.24$), and air temperature, 15-minute dose of UVB, cumulative dose of UVB, and cloud cover ($\Delta AIC_c w_i = 0.24$), respectively. Model-averaged parameter estimates that do not overlap zero (Table 2) were air temperature ($\beta = -0.215$, S.E. = 0.067, CI = -0.347 to -0.083), cumulative dose of UVB ($\beta = -0.289$, S.E. = 0.073, CI = -0.433 to -0.146), and 15-minute dose of UVB ($\beta = 0.128$, S.E. = 0.044, CI = 0.042 to 0.214). Thus, the probability of recording males in calling behavior decreased with mean air temperature and cumulative dose of UVB but increased with 15-minute dose of UVB (Fig. 3). Both the best model and the averaged model showed equally moderate predictive performance (Pearson's pseudo- $R^2 = 0.35$) and were consistent in the direction and magnitude of the effects (SM. 4).

The number of calls emitted by males was highly variable during the day (mean 5.31 ± 4.98 calls; range = 1–42), with no difference between morning (5.24 ± 0.74 sec), midday (5.39 ± 0.53 sec) and afternoon hours (4.23 ± 0.87 sec; Fig. 2B). The best-supported models to explain the number of calls emitted contained six models with ΔAIC_c weights (w_i) varying from 0.26 to 0.12, all of which contained air temperature and cumulative dose of UVB as predictors (Table 1). The best-ranked model included air temperature, luminosity, and cumulative dose of UVB ($\Delta AIC_c w_i = 0.26$). Model-averaged parameter estimates that do not overlap zero (Table 2) were air temperature ($\beta = -0.182$, S.E. = 0.043, CI = -0.267 to -0.097), and cumulative dose of UVB ($\beta = 0.107$, S.E. = 0.051, CI = 0.006 to 0.207). Thus, the




Fig. 2. Temporal patterns of the calling behavior of *Crossodactylus schmidtii* males during three periods of the day (07:00 to 18:00 h): morning hours (07:00 to 10:45 h), midday hours (11:00 to 13:45 h) and afternoon hours (14:00 to 17:45 h). Bars in (A) show the proportion of occurrence or absence of calling behavior ($n = 1936$ successive time intervals), box plots in (B) show the number of calls emitted ($n = 775$ time intervals), and in (C) show the duration of calls ($n = 775$ time intervals) emitted by *Crossodactylus schmidtii* males. The red horizontal bar inside box plots indicates the mean value, while different letters above box plots indicate significant post-hoc differences between periods by the Tukey HSD test ($p < 0.05$).

number of calls emitted by males decreased with mean air temperature but increased with cumulative dose of UVB (Fig. 4). Both the best model and the averaged model showed equally good predictive performance (Pearson's pseudo- $R^2 = 0.49$) and were consistent in the

direction and magnitude of the effects (SM. 5).

The mean duration of the advertisement calls emitted by males during the day was quite variable (mean 3.65 ± 1.16 sec, range = 0.996–9.852). However, the mean duration of calls was longer during the

Table 1. List of best-supported models ($\Delta AIC_c < 2.0$) to fit the observed data about the effect of short- and long-term cumulative doses of UVA and UVB, air temperature, light intensity and cloud cover on different proxies of calling behavior of *Crossodactylus schmidtii* males. Best-ranked models are indicated in boldface. The full model and a null model including only random terms are presented for each response variable for comparison

Response variable	Rank	Candidate model	AIC _c	ΔAIC_c	k	w _i
Occurrence of calling behavior	1	Mean air temperature + 15 min dose of UVB + Cumulative dose of UVB	2230.4	0.0	8	0.52
	2	Mean air temperature + Cumulative dose of UVA + 15 min dose of UVB + Cumulative dose of UVB	2231.9	1.5	9	0.24
	3	Cloud cover + Mean air temperature + 15 min dose of UVB + Cumulative dose of UVB	2231.9	1.6	9	0.24
	full	15 min dose of UVA * Cloud cover + 15 min dose of UVB * Cloud cover + Cumulative dose of UVA + Cumulative dose of UVB + Mean light intensity * Cloud cover + Mean air temperature * Cloud cover	2241.3	11.0	16	0.002
	null	1 + (1 site) + (1 date) + AR1(time + 0 date)	2266.9	36.5	5	< 0.001
Number of calls	1	Mean air temperature + Mean light intensity + Cumulative dose of UVB	3811.7	0.0	9	0.26
	2	Mean air temperature + Mean light intensity + Cumulative dose of UVA	3812.4	0.7	10	0.18
	3	Mean air temperature + Mean light intensity + 15 min dose of UVA	3812.6	0.9	10	0.17
	4	Mean air temperature + Mean light intensity + 15 min dose of UVA + Cumulative dose of UVA + Cumulative dose of UVB	3812.8	1.1	11	0.15
	5	Mean air temperature + 15 min dose of UVA + Cumulative dose of UVB	3813.1	1.4	9	0.13
	6	Mean air temperature + 15 min dose of UVA + Cumulative dose of UVA + Cumulative dose of UVB	3813.3	1.6	10	0.12
	full	15 min dose of UVA * Cloud cover + 15 min dose of UVB * Cloud cover + Cumulative dose of UVA + Cumulative dose of UVB + Mean light intensity * Cloud cover + Mean air temperature * Cloud cover	3821.8	10	17	0.002
	null	1 + (1 site) + (1 date) + AR1(time + 0 date)	3825.3	13.5	6	< 0.001
Mean duration of calls	1	Mean air temperature + Cumulative dose of UVB	2090.2	0.0	8	0.21
	2	Cloud cover + Mean air temperature + Cumulative dose of UVB	2090.9	0.7	9	0.15
	3	Mean air temperature + 15 min dose of UVA + Cumulative dose of UVB	2091.3	1.2	9	0.12
	4	Cloud cover * Mean air temperature + Cumulative dose of UVB	2091.3	1.2	10	0.12
	5	Mean air temperature + 15 min dose of UVB + Cumulative dose of UVB	2092.0	1.8	9	0.09
	6	Mean air temperature + 15 min dose of UVA + 15 min dose of UVB	2092.1	1.9	10	0.08
	7	Mean air temperature + Mean light intensity + Cumulative dose of UVB	2092.1	1.9	9	0.08
	8	Cloud cover + Mean air temperature + 15 min dose of UVA + Cumulative dose of UVB	2092.1	1.9	10	0.08
	9	Mean air temperature + Cumulative dose of UVA + Cumulative dose of UVB	2092.1	2.0	9	0.08
	null	1 + (1 site) + (1 date) + AR1(time + 0 date)	2096.1	6.0	6	0.01
	full	15 min dose of UVA * Cloud cover + 15 min dose of UVB * Cloud cover + Cumulative dose of UVA + Cumulative dose of UVB + Mean light intensity * Cloud cover + Mean air temperature * Cloud cover	2103.2	13.0	17	< 0.001

Note: Models are ranked by increasing order of their ΔAIC_c values. All models included the same random-effects structure: (1 | site) + (1 | date) + AR1(time | date). k = number of parameters, ΔAIC_c = difference between the AIC_c value of each model and the AIC_c value of the best-ranked model, and w_i = AIC_c weight of each model. The symbol + indicates an additive effect between variables, and the symbol * indicates an interaction between variables.




Fig. 3. Association between air temperature (A), cumulative dose of UVB (B), and 15-minutes dose of UVB (C) on the occurrence (probability) of calling behavior of *Crossodactylus schmidtii* males. The thick line represents the mean value predicted by the best-ranked model ($\Delta\text{AIC}_c w_i = 0.52$), and the gray area indicates the 95% confidence interval. Each dot represents data from successive time intervals ($n = 1936$) of passive acoustic sampling from 07:00 to 18:00 h, adjusted in the GLMMs for temporal autocorrelation. Dots are shown with a small amount of jitter for clearer visualization of superimposed data points.

Table 2. Summary of model-averaged coefficients combining the subset of best-supported models ($\Delta\text{AIC}_c < 2$) to fit the observed data about the effect of short- and long-term cumulative doses of UVA and UVB, air temperature, light intensity and cloud cover on different proxies of calling behavior of *Crossodactylus schmidtii* males. S.E. = standard error; 95% CI = 95% confidence interval

Variable	Estimate	S.E.	95% CI
Occurrence of calling behavior			
Intercept	-0.778	0.286	-1.339 to -0.218
Mean air temperature	-0.215	0.067	-0.347 to -0.083
15 min dose of UVB	0.128	0.044	0.042 to 0.214
Cumulative dose of UVB	-0.289	0.073	-0.433 to -0.146
Cumulative dose of UVA	-0.015	0.050	-0.231 to 0.107
Cloud cover	-0.007	0.026	-0.121 to 0.059
Number of calls			
Intercept	1.397	0.194	1.017 to 1.778
Mean air temperature	-0.182	0.043	-0.267 to -0.097
Mean light intensity	0.075	0.062	-0.004 to 0.202
Cumulative dose of UVB	0.107	0.051	0.006 to 0.207
Cumulative dose of UVA	-0.029	0.047	-0.164 to 0.034
15 min dose of UVA	0.028	0.037	-0.021 to 0.122
Mean duration of calls			
Intercept	3.603	0.286	3.043 to 4.164
Mean air temperature	-0.138	0.049	-0.234 to -0.043
Cumulative dose of UVB	0.104	0.052	0.001 to 0.207
Cloud cover	-0.014	0.028	-0.107 to 0.026
Cloud cover * Mean air temperature	-0.006	0.021	-0.126 to 0.027
15 min dose of UVA	0.012	0.029	-0.039 to 0.124
15 min dose of UVB	-0.006	0.022	-0.120 to 0.052
Mean light intensity	-0.002	0.018	-0.139 to 0.095
Cumulative dose of UVA	0.001	0.018	-0.101 to 0.136

Note: Estimates correspond to full model-averaged coefficients across the subset of best-supported models ($\Delta\text{AIC}_c < 2$; Table 1), with model weights incorporated in the averaging process.

morning hours (3.95 ± 0.21 sec), intermediate at midday hours (3.47 ± 0.12 sec), and lower in the afternoon hours (3.22 ± 0.13 sec; Fig. 2C). The best-supported models to explain the duration of the calls emitted by males contained nine models with ΔAIC_c weights (w_i) varying from 0.21 to 0.08, all of which contained air temperature and cumulative dose of UVB (Table 1). The best-ranked model included only air temperature

and cumulative dose of UVB as predictors (ΔAIC_c $w_i = 0.21$). Model-averaged parameter estimates that do not overlap zero (Table 2) were air temperature ($\beta = -0.138$, S.E. = 0.049, CI = -0.234 to -0.043), and cumulative dose of UVB ($\beta = 0.104$, S.E. = 0.052, CI = 0.001 to 0.207). Thus, the duration of calls emitted by males decreased with mean air temperature but increased with cumulative dose of UVB (Fig. 5). Both the best




Fig. 4. Association between air temperature (A) and cumulative dose of UVB (B) on the number of calls emitted by *Crossodactylus schmidtii* males. The thick line represents the mean value predicted by the best-ranked model (ΔAIC_c $w_i = 0.26$), and the gray area indicates the 95% confidence interval. Each dot represents data from successive time intervals ($n = 775$) of passive acoustic sampling from 07:00 to 18:00 h, adjusted in the GLMMs for temporal autocorrelation. Dots are shown with a small amount of jitter for clearer visualization of superimposed data points.




Fig. 5. Association between air Temperature (A) and cumulative dose of UVB (B) on the duration of calls emitted by *Crossodactylus schmidtii* males. The thick line represents the mean value predicted by the best-ranked model (ΔAIC_c $w_i = 0.21$), and the gray area indicates the 95% confidence interval. Each dot represents data from successive time intervals ($n = 775$) of passive acoustic sampling from 07:00 to 18:00 h, adjusted in the GLMMs for temporal autocorrelation. Dots are shown with a small amount of jitter for clearer visualization of superimposed data points.

model and the averaged models showed good predictive performance (Pearson's pseudo- $R^2 = 0.52$ and 0.53 , respectively) and were consistent in the direction and magnitude of the effects (SM. 6).

DISCUSSION

In this study we tested the association between the abiotic environment and the calling behavior of *Crossodactylus schmidtii*, a diurnal stream-breeding frog from the Neotropics. We predicted that high doses of UV reaching calling sites are associated with a reduced probability of males engaging in calling behavior. We also predicted that the investment of males in calling behavior (*i.e.*, number and duration of calls) decreases with increasing UV incidence. Overall, we expected UVB to impact male calling behavior more than UVA due to its higher DNA absorption, faster damage, and greater energy demand for repair. We found that both best-ranked and average models consistently included air temperature and long-term cumulative UVB dose as variables influencing calling behavior, with cumulative UVB effects varying in direction and magnitude across response variables. In what follows we discuss how these results support some predictions of the hypotheses tested, and we also discuss the relevance of unexpected results regarding positive associations between UVB and male calling behavior.

Higher air temperatures at calling sites were associated with a lower probability of recording males in calling behavior, as well as shorter and less frequent calls. This result agrees with well-established evidence that temperature influences anuran activity (Wells and Schwartz 2007), as daily minimum and mean temperatures often correlate positively with calling, whereas high maximum temperatures and large daily fluctuations tend to suppress it (Ospina and Villanueva-Rivera 2013; Santos et al. 2020). The association between temperature and activity reflects the basking behavior of ectotherms (Sinsch 1989; Snyder and Hammerson 1993; Centeno et al. 2015), as several diurnal species use sunlight for thermoregulation but face increasing evaporative water loss that limits activity at higher temperatures (Hatano et al. 2002; Lingnau and Bastos 2007; Ospina and Villanueva-Rivera 2013). Although this pattern is well documented, its confirmation under natural conditions is relevant in the context of concurrent solar exposure during warmer summer days, highlighting the importance of considering not only thermal but also ultraviolet components of sunlight when evaluating behavioral responses of anurans to increasing trends of environmental and climatic change.

Higher cumulative UVB doses at calling sites (but not UVA) were associated with a lower probability of recording males in calling behavior. We predicted that cumulative UVB exposure throughout the day would negatively correlate with male calling behavior, as harmful UVB doses have high genotoxic potential, leading to mutations or even cell death (Londero et al. 2019; Lundsgaard et al. 2020). At such harmful UVB doses a costly molecular mechanism is triggered to repair DNA damage (Sancar and Tang 1993; Friedberg 2003; Iyama and Wilson 2013). Since DNA damage is proportional to UVB exposure, high doses may lead to energy deficits due to the costly repair process, affecting motor, immune, and metabolic functions (Blaustein et al. 2000; Belden and Blaustein 2002; Alemanni et al. 2003; Santos et al. 2018). Although photorepair mechanisms operate on a timescale of a few femtoseconds (Kao et al. 2007), the associated energetic cost may impose a trade-off between physiological maintenance and behavioral investment (García-Huidobro et al. 2017; Pulgar et al. 2017). In anurans, calling behavior requires substantial energy, as it involves the contraction of large muscle groups relative to total body mass (Prestwich 1994; Ophir et al. 2010). Thus, under high UVB exposure, males may cease calling if they are unable to meet both the energetic demands of vocalization and the metabolic costs of DNA repair. In *C. schmidtii*, males establish and defend territories in forest streams by emitting advertisement calls throughout the daytime (Caldart et al. 2016). Consequently, ceasing calling behavior increases opportunity costs, as males failing to vocalize may lose mating opportunities and territorial control (Wells 1977; Wells and Schwartz 2007). The observed decrease in calling probability with increasing cumulative UVB exposure aligns with our prediction and suggests a direct link between harmful UVB doses and behavioral constraints in this diurnal anuran species inhabiting preserved forest streams.

The 15-minute doses of UVB (*i.e.*, a minor dose relative to cumulative UVB) were associated with an increased probability of recording males in calling behavior, and the total cumulative UVB dose incident on calling sites were associated with longer and more frequent calls. These results contradict our initial prediction that higher UVB exposure would be associated with reduced calling behavior and investment. Our data shows a pattern in which the number and duration of calls increase along the cumulative UVB gradient; however, this pattern is accompanied by a reduction in both the number of observations and data variance at higher UVB doses. This suggests that only a subset of males can sustain calling activity under high UVB exposure. This interpretation aligns with our finding that the overall

probability of calling behavior decreases as cumulative UVB exposure increases, suggesting a selective process favoring individuals with greater tolerance to UVB-induced stress. In ectothermic species, basking behavior serves not only thermoregulatory purposes but also facilitates vitamin D production (Ferguson et al. 2003) as ultraviolet B radiation plays a key role in vitamin D synthesis in vertebrates (Antwis and Browne 2009; Bouillon and Suda 2014). Vitamin D plays a critical role in calcium metabolism and muscle function, which are important for sustained calling activity. However, prolonged UVB exposure also imposes metabolic costs (Michaels and Preziosi 2013; Londero et al. 2019; Lundsgaard et al. 2020), including increased energy expenditure for DNA repair mechanisms and potential oxidative stress (Blaustein et al. 2000; Belden and Blaustein 2002; Alemanni et al. 2003; Santos et al. 2018).

Taken together, the observed pattern may suggest a potential trade-off in which males in superior physiological condition are better able to tolerate and potentially benefit from UVB exposure. Body size and energy reserves might also contribute to this pattern, since larger individuals have greater lung capacity for sustaining longer calls (Wells and Schwartz 2007; Ophir et al. 2010) and tend to possess higher lipid reserves (Simpson 1992; Scott et al. 2007). Therefore, it is plausible that males in better body condition—those with greater energy reserves and/or enhanced UVB tolerance—could be overrepresented among individuals calling under high UVB exposure. We emphasize, however, that our interpretation should be viewed with caution. Controlled manipulations of UVB exposure, combined with physiological measurements of male subjects, are needed to test this hypothesis and to evaluate whether the pattern truly reflects condition-dependent tolerance to UVB. Additionally, alternative explanations, including unmeasured environmental factors, social dynamics, or individual hormonal variation, may also contribute to the pattern observed.

CONCLUSIONS

We investigated associations between the abiotic environment and the calling behavior of *Crossodactylus schmidtii*, a diurnal frog that inhabits preserved forest streams. We found that males are less likely to engage in calling behavior during periods of higher temperatures and higher cumulative UVB doses. These results emphasize that both thermal and ultraviolet components of sunlight must be considered when evaluating how anurans respond behaviorally to ongoing environmental and climatic changes. However, contrary to our

predictions about the associations between UV exposure and calling investment, males emitted longer and more frequent calls at higher cumulative UVB doses, although fewer males did so. These findings suggest that UVB may play a role in selecting high-quality males, a possibility that warrants further experimental investigation. In addition to bringing new possibilities of research about the role of UV radiation on the calling behavior of anurans, our results showcase how heterogeneity in the abiotic environment can have contrasting behavioral effects in diurnal anurans in forest streams, organisms to which our understanding of the UV effects on behavior is still incipient.

Acknowledgments: We thank F.C. Mueller for helping in fieldwork, S. Cechin for lending audio recorders, M.C. da Rocha for assisting in obtaining the study authorization, and R. D. Schenkel for granting permission to carry out the work at Turvo State Park. We are also grateful to S. Iop for discussions on statistical analyses and to M.L.B. Santos, S. Santos, V.M. Lipinski, A.M. Tozetti, and M.T. Hartmann for suggestions that improved the manuscript. N.J. Schuch thanks CNPq Grant 300886/2016-0. This study was supported by the Conselho Nacional de Desenvolvimento Científico e Tecnológico – CNPq (Brasília, Brazil; proc. 407103/2018-0, 307063/2018-6, and 304870/2021-8) and by a CAPES/PROEX grant (proc. 23038.004173/2019-93). AAM was supported by CAPES (proc. 88887.508419/2020-00). APS is supported by CNPq (Brasília, Brazil; proc. 407103/2018-0, 307063/2018-6, 304870/2021-8, 302860/2025-8) and by CAPES/PROEX (proc. 23038.007445/2021-21, N° AUXPE: 0036/2021).

Authors' contributions: AAM, MBS, and APS conceived the study and designed the methodology; AAM and MBS developed and built the datalogger; AAM collected and organized the data with MBS under supervision of VMC; NJS provided access to cloud cover data and logistical support; VMC analyzed the data and prepared the figures; AAM wrote the first draft, and all authors revised and approved the final manuscript.

Competing interests: The authors declare no competing interests.

Availability of data and materials: All data and materials will be made available upon request.

Consent for publication: All authors consent to publication.

Ethics approval and consent to participate: This study was conducted under collection license 64137-1 issued by SEMA-RS. No individuals were captured, marked, or held in captivity, and minimal disturbance occurred due to the passive nature of data collection.

REFERENCES

- Acevedo MA, Villanueva-Rivera LJ. 2006. From the field: Using automated digital recording systems as effective tools for the monitoring of birds and amphibians. *Wildl Soc Bull* **34**:211–214. doi:10.2193/0091-7648(2006)34[211:uadrsa]2.0.co;2.
- Alemanni ME, Lozada M, Zagarese HE. 2003. Assessing sublethal effects of ultraviolet radiation in juvenile rainbow trout (*Oncorhynchus mykiss*). *Photochem Photobiol Sci* **2**:867–870. doi:10.1039/b301564e.
- Alvares CA, Stape JL, Sentelhas PC, Gonçalves JDM, Sparovek G. 2013. Köppen's climate classification map for Brazil. *Meteorol Z* **22**:711–728. doi:10.1127/0941-2948/2013/0507.
- Antwis RE, Browne RK. 2009. Ultraviolet radiation and Vitamin D₃ in amphibian health, behaviour, diet and conservation. *Comp Biochem Physiol A* **154**:184–190. doi:10.1016/j.cbpa.2009.06.008.
- Bates D, Mächler M, Bolker B, Walker S. 2015. Fitting linear mixed-effects models using lme4. *J Stat Softw* **67**:1–48. doi:10.18637/jss.v067.i01.
- Bartoń K. 2025. MuMIn: Multi-model inference. R package version 1.48.11. Available at: <https://CRAN.R-project.org/package=MuMIn> (<https://CRAN.R-project.org/package=MuMIn>). doi:10.32614/cran.package.mumin.
- Belden LK, Blaustein AR. 2002. Exposure of red-legged frog embryos to ambient UV-B radiation in the field negatively affects larval growth and development. *Oecologia* **130**:551–554. doi:10.1007/s00442-001-0843-y.
- Blaustein AR, Belden LK. 2003. Amphibian defenses against ultraviolet-B radiation. *Evol Dev* **5**:89–97. doi:10.1046/j.1525-142x.2003.03014.x.
- Blaustein AR, Chivers DP, Kats LB, Kiesecker JM. 2000. Effects of ultraviolet radiation on locomotion and orientation in roughskin newts (*Taricha granulosa*). *Ethology* **106**:227–234. doi:10.1046/j.1439-0310.2000.00519.x.
- Blumstein DT, Mennill DJ, Clemins P, Girod L, Yao K et al. 2011. Acoustic monitoring in terrestrial environments using microphone arrays: applications, technological considerations and prospectus. *J Appl Ecol* **48**:758–767. doi:10.1111/j.1365-2664.2011.01993.x.
- Bouillon R, Suda T. 2014. Vitamin D: calcium and bone homeostasis during evolution. *Bonekey Rep* **3**:480. doi:10.1038/bonekey.2013.214.
- Brenner M, Hearing VJ. 2008. The protective role of melanin against UV damage in human skin. *Photochem Photobiol* **84**:539–549. doi:10.1111/j.1751-1097.2007.00226.x.
- Brooks ME, Kristensen K, Van Benthem KJ, Magnusson A, Berg CW et al. 2017. glmmTMB balances speed and flexibility among packages for zero-inflated generalized linear mixed modeling. *The R Journal* **9**:378–400. doi:10.32614/rj-2017-066.
- Caldart VM, dos Santos MB, Machado G. 2022. Function of a multimodal signal: a multiple hypothesis test using a robot frog. *J Anim Ecol* **91**:701–714. doi:10.1111/1365-2656.13620.
- Caldart VM, Iop S, Cechin SZ. 2011. Vocalizations of *Crossodactylus schmidtii* Gallardo, 1961 (Anura, Hylodidae): advertisement call and aggressive call. *North-West J Zool* **7**:118–124.
- Caldart VM, Iop S, Cechin SZ. 2014. Social interactions in a neotropical stream frog reveal a complex repertoire of visual signals and the use of multimodal communication. *Behaviour* **151**:719–739. doi:10.1163/1568539x-00003165.
- Caldart VM, Iop S, Lingnau R, Cechin SZ. 2016. Calling activity of a stream-breeding frog from the austral neotropics: temporal patterns of activity and the role of environmental factors. *Herpetologica* **72**:90–97. doi:10.1655/herpetologica-d-15-00029.
- Centeno FC, Antoniazzi MM, Andrade DV, Kodama RT, Sciani JM et al. 2015. Anuran skin and basking behavior: the case of the treefrog *Bokermannohyla alvarengai* (Bokermann, 1956). *J Morphol* **276**:1172–1182. doi:10.1002/jmor.20407.
- Cockell C, Blaustein AR (eds). 2001. Ecosystems, evolution, and ultraviolet radiation. Springer New York, NY. doi:10.1007/978-1-4757-3486-7.
- Cortat B, Garcia CCM, Quinet A, Schuch AP, de Lima-Bessa KM et al. 2013. The relative roles of DNA damage induced by UVA irradiation in human cells. *Photochem Photobiol Sci* **12**:1483–1495. doi:10.1039/c3pp50023c.
- Daan S. 1981. Adaptive daily strategies in behavior. In: Aschoff J (ed) *Biological rhythms*. Springer, pp. 275–298. doi:10.1007/978-1-4615-6552-9_15.
- Davison AC, Hinkley DV. 1997. Bootstrap methods and their application. Cambridge University Press. doi:10.1017/cbo9780511802843.
- Dong Q, Svoboda K, Tiersch TR, Monroe WT. 2007. Photobiological effects of UVA and UVB light in zebrafish embryos: evidence for a competent photorepair system. *J Photochem Photobiol B* **88**:137–146. doi:10.1016/j.jphotobiol.2007.07.002.
- Ferguson GW, Gehrman WH, Karsten KB, Hammack SH, McRae M et al. 2003. Do panther chameleons bask to regulate endogenous vitamin D₃ production? *Physiol Biochem Zool* **76**:52–59. doi:10.1086/374276.
- Frederick JE, Snell HE, Haywood EK. 1989. Solar ultraviolet radiation at the earth's surface. *Photochem Photobiol* **50**:443–450. doi:10.1111/j.1751-1097.1989.tb05548.x.
- Friedberg EC. 2003. DNA damage and repair. *Nature* **421**:436–440. doi:10.1038/nature01408.
- García-Huidobro MR, Aldana M, Duarte C, Galbán-Malagón C, Pulgar J. 2017. Seawater-temperature and UV-radiation interaction modifies oxygen consumption, digestive process and growth of an intertidal fish. *Mar Environ Res* **129**:408–412. doi:10.1016/j.marenvres.2017.06.013.
- Hamzavi I, Lui H. 2005. Using light in dermatology: an update on lasers, ultraviolet phototherapy, and photodynamic therapy. *Dermatol Clin* **23**:199–207. doi:10.1016/j.det.2004.11.001.
- Hartig F. 2022. DHARMA: residual diagnostics for hierarchical (multi-level/mixed) regression models. R package. doi:10.32614/cran.package.dharma.
- Hatano FH, Rocha CFD, Van Sluys M. 2002. Environmental factors affecting calling activity of a tropical diurnal frog (*Hylodes phyllodes*). *J Herpetol* **36**:314–318. doi:10.1670/0022-1511(2002)036[0314:efacao]2.0.co;2.
- Hohman EE, Martin BR, Lachcik PJ, Gordon DT, Fleet JC et al. 2011. Bioavailability and efficacy of vitamin D₂ from UV-irradiated yeast in growing, vitamin D-deficient rats. *J Agric Food Chem* **59**:2341–2346. doi:10.1021/jf104679c.
- Iyama T, Wilson DM. 2013. DNA repair mechanisms in dividing and non-dividing cells. *DNA Repair* **12**:620–636. doi:10.1016/j.dnarep.2013.04.015.
- James G, Witten D, Hastie T, Tibshirani R. 2013. An introduction to statistical learning. Springer New York, NY. doi:10.1007/978-1-4614-7138-7_2.
- Kao YT, Saxena C, Wang L, Sancar A, Zhong D. 2007. Femtochemistry in enzyme catalysis: DNA photolyase. *Cell Biochem Biophys* **48**:32–44. doi:10.1007/s12013-007-0034-5.

- Kats LB, Bucciarelli GM, Schlais DE, Blaustein AR, Han BA. 2012. Ultraviolet radiation influences perch selection by a Neotropical poison-dart frog. *PLoS ONE* 7:e51364. doi:10.1371/journal.pone.0051364.
- Kats LB, Kiesecker JM, Chivers DP, Blaustein AR. 2000. Effects of UV-B radiation on anti-predator behavior in three species of amphibians. *Ethology* 106:921–931. doi:10.1046/j.1439-0310.2000.00608.x.
- Kienzler A, Tronchère X, Devaux A, Bony S. 2013. UV-induced nucleotide excision repair (NER) and photoreactivation repair (PER) in two trout fish cell lines used in ecotoxicological assessment studies. *J Photochem Photobiol B* 125:51–55. doi:10.1016/j.jphotobiol.2013.05.004.
- Kouwenberg JHM, Browman HI, Cullen JJ, Davis RF, St-Pierre JF et al. 1999. Biological weighting of ultraviolet (280–400 nm) induced mortality in marine zooplankton and fish. I. Atlantic cod (*Gadus morhua*) eggs. *Mar Biol* 134:269–284. doi:10.1007/s002270050545.
- Lamare MD, Barker MF, Lesser MP, Marshall C. 2006. DNA photorepair in echinoid embryos: effects of temperature on repair rate in Antarctic and non-Antarctic species. *J Exp Biol* 209:5017–5028. doi:10.1242/jeb.02598.
- Larsen AS, Schmidt JH, Stapleton H, Kristenson H, Betchkal D et al. 2021. Monitoring the phenology of the wood frog breeding season using bioacoustic methods. *Ecol Indic* 131:108142. doi:10.1016/j.ecolind.2021.108142.
- Lenth R, Piaskowski J. 2025. emmeans-package: Estimated marginal means, aka least-squares means. R package version 2.0.0. doi:10.32614/cran.package.emmeans.
- Lesser MP, Barry TM. 2003. Survivorship, development, and DNA damage in echinoderm embryos and larvae exposed to ultraviolet radiation (290–400 nm). *J Exp Mar Biol Ecol* 292:75–91. doi:10.1016/s0022-0981(03)00141-2.
- Lingnau R, Bastos RP. 2007. Vocalizations of the Brazilian torrent frog *Hylodes heyeri* (Anura: Hylodidae): repertoire and influence of air temperature on advertisement call variation. *J Nat Hist* 41:1227–1235. doi:10.1080/00222930701395626.
- Londero JEL, Santos CP, Segatto ALA, Schuch AP. 2017. Impacts of UVB radiation on food consumption of forest specialist tadpoles. *Ecotoxicol Environ Saf* 143:12–18. doi:10.1016/j.ecoenv.2017.05.002.
- Londero JEL, Santos MB, Schuch AP. 2019. Impact of solar UV radiation on amphibians: focus on genotoxic stress. *Mutat Res Genet Toxicol Environ Mutagen* 842:14–21. doi:10.1016/j.mrgentox.2019.03.003.
- Lüdtke D, Ben-Shachar MS, Patil I, Waggoner P, Makowski D. 2021. performance: An R package for assessment, comparison and testing of statistical models. *J Open Source Softw* 6:3139. doi:10.21105/joss.03139.
- Lundsgaard NU, Cramp RL, Franklin CE. 2020. Effects of ultraviolet-B radiation on physiology, immune function and survival is dependent on temperature: implications for amphibian declines. *Conserv Physiol* 8:coaa002. doi:10.1093/conphys/coaa002.
- Maluf JRT. 2000. Nova classificação climática do Estado do Rio Grande do Sul. *Rev Bras Agrometeorol* 8:141–150.
- Markovitsi D, Gustavsson T, Banyasz A. 2010. Absorption of UV radiation by DNA: spatial and temporal features. *Mutat Res Mutat Res* 704:21–28. doi:10.1016/j.mrrev.2009.11.003.
- Martins ÍM, Vasconcellos ADS, Mota T, Eterovick PC. 2021. Detectability is in the eye of the beholder—the role of UV reflectance on tadpole detection and predation by a passerine bird. *Behav Ecol Sociobiol* 75:1–14. doi:10.1007/s00265-021-02983-9.
- Michaels CJ, Preziosi RF. 2013. Basking behaviour and ultraviolet B radiation exposure in a wild population of *Pelophylax lessonae* in northern Italy. *Herpetol Bull* 124:1–8.
- Nakagawa S, Schielzeth H. 2013. A general and simple method for obtaining R^2 from generalized linear mixed-effects models. *Methods Ecol Evol* 4:133–142. doi:10.1111/j.2041-210x.2012.00261.x.
- Neer RM. 1975. The evolutionary significance of vitamin D, skin pigment, and ultraviolet light. *Am J Phys Anthropol* 43:409–416. doi:10.1002/ajpa.1330430322.
- Ophir AG, Schrader SB, Gillooly JF. 2010. Energetic cost of calling: general constraints and species-specific differences. *J Evol Biol* 23:1564–1569. doi:10.1111/j.1420-9101.2010.02005.x.
- Ospina OE, Villanueva-Rivera LJ, Corrada-Bravo CJ, Aide TM. 2013. Variable response of anuran calling activity to daily precipitation and temperature: implications for climate change. *Ecosphere* 4:1–12. doi:10.1890/es12-00258.1.
- Prestwich KN. 1994. The energetics of acoustic signaling in anurans and insects. *Am Zool* 34:625–643. doi:10.1093/icb/34.6.625.
- Pulgar J, Waldisparg M, Galbán-Malagón C, Maturana D, Pulgar VM et al. 2017. UV radiation impacts body weight, oxygen consumption, and shelter selection in the intertidal vertebrate *Girella laevis*. *Sci Total Environ* 578:317–322. doi:10.1016/j.scitotenv.2016.10.157.
- R Core Team. 2022. R: a language and environment for statistical computing. R Foundation for Statistical Computing, Vienna. doi:10.32614/r.manuals.
- Sancar A, Tang MS. 1993. Nucleotide excision repair. *Photochem Photobiol* 57:905–921. doi:10.1111/j.1751-1097.1993.tb09233.x.
- Santos CP, Londero JEL, Santos MB, Santos Feltrin R, Loebens L et al. 2018. Sunlight-induced genotoxicity and damage in keratin structures decrease tadpole performance. *J Photochem Photobiol B* 181:134–142. doi:10.1016/j.jphotobiol.2018.03.013.
- Santos MBD, Madalozzo B, Tozetti AM, Loebens L, Cechin SZ. 2020. Climatic dependence in the daily and seasonal calling activity of anurans from coastal wetlands of southernmost Brazil. *J Nat Hist* 54:2037–2052. doi:10.1080/00222933.2020.1833998.
- Schielzeth H. 2010. Simple means to improve the interpretability of regression coefficients. *Methods Ecol Evol* 1:103–113. doi:10.1111/j.2041-210x.2010.00012.x.
- Schuch AP, Menck CFM. 2010. The genotoxic effects of DNA lesions induced by artificial UV-radiation and sunlight. *J Photochem Photobiol B* 99:111–116. doi:10.1016/j.jphotobiol.2010.03.004.
- Schuch AP, Santos MB, Lipinski VM, Peres LV, DSantos et al. 2015. Identification of influential events concerning the Antarctic ozone hole over southern Brazil and the biological effects induced by UVB and UVA radiation in an endemic treefrog species. *Ecotoxicol Environ Saf* 118:190–198. doi:10.1016/j.ecoenv.2015.04.029.
- Schuch AP, Moreno NC, Schuch NJ, Menck CFM, Garcia CCM. 2017. Sunlight damage to cellular DNA: focus on oxidatively generated lesions. *Free Radic Biol Med* 107:110–124. doi:10.1016/j.freeradbiomed.2017.01.029.
- Scott DE, Casey ED, Donovan MF, Lynch TK. 2007. Amphibian lipid levels at metamorphosis correlate to post-metamorphic terrestrial survival. *Oecologia* 153:521–532. doi:10.1007/s00442-007-0755-6.
- SEMA. 2005. Plano de manejo do Parque Estadual do Turvo. Secretaria Estadual do Meio Ambiente.
- Simpson AL. 1992. Differences in body size and lipid reserves between maturing and nonmaturing Atlantic salmon parr, *Salmo salar* L. *Can J Zool* 70:1737–1742. doi:10.1139/z92-241.
- Sinsch U. 1989. Behavioural thermoregulation of the Andean toad (*Bufo spinulosus*) at high altitudes. *Oecologia* 80:32–38. doi:10.1007/bf00789928.
- Snyder GK, Hammerson GA. 1993. Interrelationships between water

- economy and thermo-regulation in the canyon tree-frog *Hyla arenicolor*. *J Arid Environ* **25**:321–329. doi:10.1006/jare.1993.1065.
- Souza J, Macedo AS, Martins FR, Pereira EB. 2013. Metodologias de estimativa de cobertura de nuvens e seus resultados no modelo BRASIL-SR. *In: Anais do XVI Simpósio Brasileiro de Sensoriamento Remoto, Foz do Iguaçu, Brazil*, pp. 6207–6213. doi:10.5967/cbens.2018.675.
- Symonds MRE, Moussalli A. 2011. A brief guide to model selection, multimodel inference and model averaging in behavioural ecology using Akaike's information criterion. *Behav Ecol Sociobiol* **65**:13–21. doi:10.1007/s00265-010-1037-6.
- Van Sluys M, Guido-Castro P. 2011. Influence of temperature and photoperiod on the activity of *Melanophryniscus moreirae* (Anura: Bufonidae) on the Itatiaia Plateau, southeastern Brazil. *South Am J Herpetol* **6**:43–48. doi:10.2994/057.006.0106.
- Wells KD. 1977. The social behaviour of anuran amphibians. *Anim Behav* **25**:666–693. doi:10.1016/0003-3472(77)90118-x.
- Wells KD, Schwartz JJ. 2007. The behavioral ecology of anuran communication. *In: Narins PM et al. (eds) Hearing and sound communication in amphibians*. Springer, pp. 44–86. doi:10.1007/978-0-387-47796-1_3.
- Xue Y, Fleet JC. 2009. Intestinal vitamin D receptor is required for normal calcium and bone metabolism in mice. *Gastroenterology* **136**:1317–1327. doi:10.1053/j.gastro.2008.12.051.
- Yagura T, Schuch AP, Garcia CCM, Rocha CRR, Moreno NC et al. 2017. Direct participation of DNA in the formation of singlet oxygen and base damage under UVA irradiation. *Free Radic Biol Med* **108**:86–93. doi:10.1016/j.freeradbiomed.2017.03.018.
- Zagarese HE, Williamson CE. 2001. The implications of solar UV radiation exposure for fish and fisheries. *Fish Fish* **2**:250–260. doi:10.1046/j.1467-2960.2001.00048.x.
- ## Supplemental materials
- SM. 1.** Validation test of the UV data logger. (download)
- SM. 2.** Patterns of UVA and UVB incidence in control and sampling sites. (download)
- SM. 3.** Specifications of the Generalized Linear Mixed Models used to test the hypotheses outlined in the main text. (download)
- SM. 4.** Results and diagnostics of models following the four-step statistical analysis procedure for the dependent variable 'Occurrence of calling behavior'. (download)
- SM. 5.** Results and diagnostics of models following the four-step statistical analysis procedure for the dependent variable 'Number of calls emitted'. (download)
- SM. 6.** Results and diagnostics of models following the four-step statistical analysis procedure for the dependent variable 'Mean duration of calls'. (download)